

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number  
**WO 03/029215 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 211/70**,  
401/06, 407/06, 409/06, 417/06, A61K 31/4523, 31/445,  
A61P 25/04

Montreal, 7171 Frederick-Banting, St. Laurent, Montreal,  
Québec H4S 1Z9 (CA). **WEI, Zhongyong** [CA/CA]; As-  
traZeneca R & D Montreal, 7171 Frederick-Banting, St.  
Laurent, Montreal, Québec H4S 1Z9 (CA).

(21) International Application Number: PCT/SE02/01804

(74) Agent: **GLOBAL INTELLECTUAL PROPERTY**; As-  
traZeneca AB, S-151 85 Södertälje (SE).

(22) International Filing Date: 2 October 2002 (02.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0103313-3 3 October 2001 (03.10.2001) SE

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **AS-  
TRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).


(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BROWN, William**  
[CA/CA]; AstraZeneca R & D Montreal, 7171 Frederick-  
Banting, St. Laurent, Montreal, Québec H4S 1Z9 (CA).  
**WALPOLE, Christopher** [GB/CA]; AstraZeneca R & D

[Continued on next page]

(54) Title: 4[PIPERIDIN-4-YLIDEN-(3-CARBAMOYLPHENYL)METHYL] BENZAMIDE DERIVATIVES AND THEIR USE FOR THE TREATMENT OF PAIN, SPINAL INJURIES OR GASTROINTESTINAL DISORDERS



(57) Abstract: Compounds of general formula (I) wherein R<sup>1</sup> is selected from any one of phenyl, pyridinyl, thienyl, furanyl, imidazolyl, pyrrolyl and triazolyl; where each R<sup>1</sup> phenyl ring and R<sup>1</sup> heteroaromatic ring may optionally and independently be further substituted by 1, 2 or 3 substituents selected from straight and branched C<sub>1</sub>-C<sub>6</sub> alkyl, NO<sub>2</sub>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, chloro, fluoro, bromo, and iodo. The substitutions on the phenyl ring and on the heteroaromatic ring may take place in any position on said ring systems; are disclosed and claimed in the present application, as well as their pharmaceutically acceptable salts and pharmaceutical compositions comprising the novel compounds and their use in therapy, in particular in the management of pain.

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*

- *of inventorship (Rule 4.17(iv)) for US only*

**Published:**

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

4[piperidin-4-yliden-(3-carbamoylphenyl)methyl] benzamide derivatives and their use for the treatment of pain, spinal injuries or gastrointestinal disorders

### **Field of the Invention**

The present invention is directed to novel compounds, to a process for their preparation,  
5 their use and pharmaceutical compositions comprising the novel compounds. The novel compounds are useful in therapy, and in particular for the treatment of pain.

### **Background of the Invention**

The  $\delta$  receptor has been identified as having a role in many bodily functions such as circulatory and pain systems. Ligands for the  $\delta$  receptor may therefore find potential use as  
10 analgesics, and/or as antihypertensive agents. Ligands for the  $\delta$  receptor have also been shown to possess immunomodulatory activities.

The identification of at least three different populations of opioid receptors ( $\mu$ ,  $\delta$  and  $\kappa$ ) is now well established and all three are apparent in both central and peripheral nervous systems of many species including man. Analgesia has been observed in various animal models when one or  
15 more of these receptors has been activated.

With few exceptions, currently available selective opioid  $\delta$  ligands are peptidic in nature and are unsuitable for administration by systemic routes. One example of a non-peptidic  $\delta$ -agonist is SNC80 (*Bilsky E.J. et al., Journal of Pharmacology and Experimental Therapeutics*, 273(1), pp. 359-366 (1995)). There is however still a need for selective  $\delta$ -agonists having not only improved  
20 selectivity, but also an improved side-effect profile.


Thus, the problem underlying the present invention was to find new analgesics having improved analgesic effects, but also with an improved side-effect profile over current  $\mu$  agonists, as well as having improved systemic efficacy.

Analgesics that have been identified and exist in the prior art have many disadvantages in that  
25 they suffer from poor pharmacokinetics and are not analgesic when administered by systemic routes. Also, it has been documented that preferred  $\delta$  agonist compounds, described within the prior art, show significant convulsive effects when administered systemically.


We have now found that certain compounds not specifically disclosed by, but included within the scope of WO 98/28275, exhibit surprisingly improved  $\delta$ -agonist properties and in vivo potency.


### Outline of the invention

5 The novel compounds according to the present invention are defined by the formula I wherein








10  $R^1$  is selected from any one of

(i) phenyl; 

(ii) pyridinyl  ;

(iii) thienyl  ;

(iv) furanyl  ;






where each  $R^1$  phenyl ring and  $R^1$  heteroaromatic ring may optionally and independently be further substituted by 1, 2 or 3 substituents independently selected from straight and branched  $C_1$ - $C_6$  alkyl,  $NO_2$ ,  $CF_3$ ,  $C_1$ - $C_6$  alkoxy, chloro, fluoro, bromo, and iodo. The substitutions on the phenyl ring and on the heteroaromatic ring may take place in any position on said ring systems.


Particularly, novel compounds according to the present invention are defined by the formula

I

wherein  $R^1$  is selected from any one of



(iv) furanyl



Within the scope of the invention are also salts and enantiomers of the compounds of the formula I.

When the phenyl ring and the heteroaromatic ring(s) are substituted, the preferred substituents are independently selected from any one of CF<sub>3</sub>, methyl, iodo, bromo, fluoro and chloro.

The novel compounds of the present invention are useful in therapy, especially for the treatment of various pain conditions such as chronic pain, neuropathic pain, acute pain, cancer pain, pain caused by rheumatoid arthritis, migraine, visceral pain etc. This list should however not be interpreted as exhaustive.

Compounds of the invention are useful as immunomodulators, especially for autoimmune diseases, such as arthritis, for skin grafts, organ transplants and similar surgical needs, for collagen diseases, various allergies, for use as anti-tumour agents and anti viral agents.

Compounds of the invention are useful in disease states where degeneration or dysfunction of opioid receptors is present or implicated in that paradigm. This may involve the use of isotopically labelled versions of the compounds of the invention in diagnostic techniques and imaging applications such as positron emission tomography (PET).

Compounds of the invention are useful for the treatment of diarrhoea, depression, anxiety, urinary incontinence, various mental illnesses, cough, lung oedema, various gastro-intestinal disorders, spinal injury and drug addiction, including the treatment of alcohol, nicotine, opioid and other drug abuse and for disorders of the sympathetic nervous system for example hypertension.


Compounds of the invention are useful as an analgesic agent for use during general anaesthesia and monitored anaesthesia care. Combinations of agents with different properties are often used to achieve a balance of effects needed to maintain the anaesthetic state (e.g. amnesia, analgesia, muscle relaxation and sedation). Included in this combination are inhaled anaesthetics, hypnotics, anxiolytics, neuromuscular blockers and opioids.

Also within the scope of the invention is the use of any of the compounds according to the

formula I above, for the manufacture of a medicament for the treatment of any of the conditions discussed above.

A further aspect of the invention is a method for the treatment of a subject suffering from any of the conditions discussed above, whereby an effective amount of a compound according to the formula I above, is administered to a patient in need of such treatment.

A further aspect of the present invention is intermediates of the general formula II




wherein PG is a urethane protecting group such as Boc or CBZ, or a benzyl or substituted benzyl protecting group, such as 2,4-dimethoxybenzyl.

#### Methods of preparation

#### EXAMPLES

The invention will now be described in more detail by the following Schemes and Examples, which are not to be construed as limiting the invention.



Scheme 1: Synthetic Route to Compounds of the Present Invention

**N,N-Diethyl-4-[N-Boc-piperidin-4-ylidene(3-carboxyphenyl)-methyl]-benzamide (2a).**

A mixture of 4-[bromo-(4-diethylcarbamoyl-phenyl)-methylene]-piperidine-1-carboxylic acid tert-butyl ester (**1a**, 451 mg, 1.0 mmol), 3-carboxyphenyl boronic acid (330 mg, 2.0 mmol), 2M Na<sub>2</sub>CO<sub>3</sub> (3 mL), and tetrakis(triphenyl phosphine) palladium(0) (25 mg) in toluene (degassed, 10 mL) and ethanol (degassed, 10 mL) was refluxed at 90 °C for 4 hrs under N<sub>2</sub>. The reaction mixture was then quenched with aqueous NH<sub>4</sub>Cl after cooling down to 0 °C, and extracted with ethyl acetate (2 x 50 mL). The combined organic phases were washed with brine, dried over MgSO<sub>4</sub>, and evaporated to give a crude product, which was purified by flash silica gel column to provide the desired compound **2a** (345 mg, 70 %): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.15 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 1.23 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 1.47 (9 H, s, C(CH<sub>3</sub>)<sub>3</sub>), 2.31 (2 H, m, piperidine CH<sub>2</sub>-), 2.35 (2 H, m,



piperidine  $\text{CH}_2$ -), 3.30 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.48 (4 H, m, piperidine  $\text{CH}_2$ -), 3.54 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 7.14 (2 H, d,  $J = 8.0$  Hz, ArH), 7.24 (1 H, m, ArH), 7.33 (2 H, d,  $J = 8.0$  Hz, ArH), 7.42 (1 H, t,  $J = 7.6$  Hz, ArH), 7.86 (1 H, s, ArH), 7.97 (1 H, d,  $J = 7.6$  Hz, ArH). IR (NaCl) 2976, 1718, 1691, 1598, 1430, 1233, 1166  $\text{cm}^{-1}$ .

5 **N,N-Diethyl-4-[1-(benzyl-piperidin-4-ylidene-(3-carboxyphenyl)-methyl)-benzamide (2b).**

Method as for **2a** using **1b** (441 mg, 1.0 mmol) and 3-carboxyphenyl boronic acid (330 mg, 2.0 mmol) provided **2b** (325 mg, 67 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.11 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.22 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 2.63 (4 H, m, piperidine  $\text{CH}_2$ -), 2.90 (4 H, m, piperidine  $\text{CH}_2$ -), 3.26  
10 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.52 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 4.00 (2 H, s,  $\text{CH}_2\text{N}$ -), 7.11 (2 H, d,  $J = 8.0$  Hz, ArH), 7.16 (1H, m, ArH), 7.30 (6 H, m, ArH), 7.42 (2 H, m, ArH), 7.84 (1 H, s, ArH), 7.96 (1 H, d,  $J = 7.6$  Hz, ArH).

**N,N-Diethyl-4-[1-benzylpiperidin-4-ylidene(3-carbamoylphenyl)-methyl]-benzamide (3b).**

15 241 mg (0.5 mmol) of N,N-diethyl-4-[piperidin-4-ylidene(3-carboxyphenyl)-methyl]-benzamide (**2b**), 780 mg (1.5 mmol) of PyBOP, and 200 mg (1.5 mmol) of HOBt were dissolved in 4 mL DMF. 0.58 mL (4 mmol) of DIPEA and 50 mg (10.0 mmol) of  $\text{NH}_4\text{Cl}$  were added successively. After stirred for 0.5 h at room temperature, the reaction mixture was quenched with water and ethyl ether. The white precipitates were collected as the desired product (**3b**, 152 mg, 63  
20 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.10 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.21 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 2.62 (4 H, m, piperidine  $\text{CH}_2$ -), 2.89 (4 H, m, piperidine  $\text{CH}_2$ -), 3.26 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.52 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 4.00 (2 H, s,  $\text{CH}_2\text{N}$ -), 7.11 (2 H, d,  $J = 8.0$  Hz, ArH), 7.16 (1H, m, ArH), 7.30 (6 H, m, ArH), 7.42 (2 H, m, ArH), 7.84 (1 H, s, ArH), 7.96 (1 H, d,  $J = 7.6$  Hz, ArH);  
*Anal. Calcd. for*  $\text{C}_{31}\text{H}_{35}\text{N}_3\text{O}_2 \cdot 6.0 \text{ HCl}$ : C, 53.16 %; H, 5.90 %; *Found*: C, 53.07 %; H, 5.54  
25 %.

**N,N-Diethyl-4-[piperidin-4-ylidene(3-carbamoylphenyl)-methyl]-benzamide (4).**

Method as for **3b** using **2a** (150 mg, 0.30 mmol) provided **3a** (105 mg, 70 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.14 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.22 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.46 (9 H, s,  $\text{C}(\text{CH}_3)_3$ ), 2.28 (2 H, m, piperidine  $\text{CH}_2$ -), 2.33 (2 H, m, piperidine  $\text{CH}_2$ -), 3.30 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.46

(4 H, m, piperidine CH-), 3.55 (2 H, br m, CH<sub>3</sub>CH<sub>2</sub>N-), 7.13 (2 H, d, J = 8.0 Hz, ArH), 7.30 (3 H, m, ArH), 7.39 (1 H, t, J = 7.6 Hz, ArH), 7.61 (1 H, s, ArH), 7.68 (1 H, d, J = 7.6 Hz, ArH).

The above product (**3a**) was treated with 4.0 M HCl in dioxane (10 mL) at room temperature for 4 h. After evaporation, the residue was dissolved in H<sub>2</sub>O (10 mL) and impurities were extracted with ethyl acetate (2 x 20 mL). The aqueous phase was basified with NH<sub>4</sub>OH and extracted with ethyl acetate (3 x 20 mL). The combined organic phases were washed with brine, dried over MgSO<sub>4</sub> and evaporated to give **4** in quantitative yield: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.15 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 1.23 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 1.90 (2 H, br, NH<sub>2</sub>), 2.31 (2 H, m, piperidine CH-), 2.34 (2 H, m, piperidine CH-), 2.92 (4 H, m, piperidine CH-), 3.32 (2 H, br m, CH<sub>3</sub>CH<sub>2</sub>N-), 3.55 (2 H, br m, CH<sub>3</sub>CH<sub>2</sub>N-), 7.13 (2 H, d, J = 8.0 Hz, ArH), 7.30 (3 H, m, ArH), 7.38 (1 H, t, J = 7.6 Hz, ArH), 7.58 (1 H, s, ArH), 7.66 (1 H, d, J = 7.6 Hz, ArH). IR (NaCl) 3307, 2973, 1668, 1615, 1435, 1383, 1289 cm<sup>-1</sup>; *Anal. Calcd. for* C<sub>24</sub>H<sub>29</sub>N<sub>3</sub>O<sub>2</sub> 2.8 HCl: C, 58.40 %; H, 6.49 %; *Found*: C, 58.46 %; H, 6.57 %.

**N,N-Diethyl-4-[1-(2-thiophene)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5a).**

To a mixture of N,N-diethyl-4-[piperidin-4-ylidene(3-carbamoylphenyl)-methyl]-benzamide (**4**, 196 mg, 0.5 mmol), 2-thiophenecarboxaldehyde (112 mg, 1.0 mmol), acetic acid (0.1 mL) in MeOH (10 mL) was added NaBH<sub>3</sub>(CN) (200 mg) in portions. The reaction mixture was stirred for 4 h at room temperature, and then quenched with aqueous NH<sub>4</sub>Cl, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 50 mL). The combined organic phases were washed with brine, dried over MgSO<sub>4</sub>, and evaporated to give a crude product, which was purified by flash silica gel column to give the desired product (**5a**, 216 mg, 89 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.13 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 1.22 (3 H, br m, CH<sub>3</sub>CH<sub>2</sub>-), 2.36 (2 H, m, piperidine CH-), 2.42 (2 H, m, piperidine CH-), 2.54 (4 H, m, piperidine CH-), 3.26 (2 H, br m, CH<sub>3</sub>CH<sub>2</sub>N-), 3.52 (2 H, br m, CH<sub>3</sub>CH<sub>2</sub>N-), 3.76 (2 H, s, CH<sub>2</sub>N-), 5.60 (1 H, br, NH), 6.06 (1 H, br, NH), 6.91 (1 H, s, ArH), 6.94 (1 H, m, ArH), 7.12 (2 H, d, J = 8.0 Hz, ArH), 7.26 (1 H, m, ArH), 7.30 (3 H, m, ArH), 7.37 (1 H, t, J = 7.6 Hz, ArH), 7.56 (1 H, s, ArH), 7.64 (1 H, d, J = 7.6 Hz, ArH). IR (NaCl) 3352, 2973, 1668, 1614, 1434, 1290 cm<sup>-1</sup>; *Anal. Calcd. for* C<sub>29</sub>H<sub>33</sub>N<sub>3</sub>O<sub>2</sub>S 2.3 HCl: C, 60.95 %; H, 6.23 %; *Found*: C, 60.96 %; H, 6.42 %.

**N,N-Diethyl-4-[1-(2-furfuryl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl)-benzamide (5b).**

Method as for **5a** using **4** (196 mg, 0.5 mmol) and 2-furaldehyde (96 mg, 1.0 mmol) provided **5b** (158 mg, 67 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.12 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.22 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 2.41 (4 H, m, piperidine  $\text{CH}$ -), 2.83 (4 H, m, piperidine  $\text{CH}$ -), 3.27 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.52 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.88 (2 H, s,  $\text{CH}_2\text{N}$ -), 6.18 (1 H, br, NH), 6.35 (1 H, m, ArH), 6.42 (1 H, m, ArH), 6.86 (1 H, m, NH), 7.12 (2 H, d,  $J = 8.0$  Hz, ArH), 7.27 (3 H, m, ArH), 7.32 (1 H, m, ArH), 7.41 (1 H, s, ArH), 7.61 (1 H, s, ArH), 7.69 (1 H, d,  $J = 7.6$  Hz, ArH); *Anal. Calcd. for*  $\text{C}_{29}\text{H}_{33}\text{N}_3\text{O}_3 \cdot 3.0 \text{ HCl}$ : C, 59.95 %; H, 6.25 %; *Found*: C, 59.68 %; H, 5.98 %.

**N,N-Diethyl-4-[1-(3-furfuryl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl)-benzamide (5c).**

Method as for **5a** using **4** (196 mg, 0.5 mmol) and 3-furaldehyde (96 mg, 1.0 mmol) provided **5c** (143 mg, 61 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.13 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.25 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 2.42 (4 H, m, piperidine  $\text{CH}$ -), 2.70 (4 H, m, piperidine  $\text{CH}$ -), 3.26 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.52 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.62 (2 H, s,  $\text{CH}_2\text{N}$ -), 5.80 (1 H, br, NH), 6.42 (1 H, br, NH), 6.46 (1 H, s, ArH), 7.12 (2 H, d,  $J = 8.0$  Hz, ArH), 7.28 (3 H, m, ArH), 7.36 (1 H, t,  $J = 7.6$  Hz, ArH), 7.42 (2 H, m, ArH), 7.59 (1 H, s, ArH), 7.66 (1 H, d,  $J = 7.6$  Hz, ArH); *Anal. Calcd. for*  $\text{C}_{29}\text{H}_{33}\text{N}_3\text{O}_3 \cdot 3.1 \text{ HCl}$ : C, 59.58 %; H, 6.22 %; *Found*: C, 59.44 %; H, 6.45 %.

**N,N-Diethyl-4-[1-(2-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5d).**

Method as for **5a** using **4** (196 mg, 0.5 mmol) and 2-pyridinecarboxaldehyde (107 mg, 1.0 mmol) provided **5d** (35 mg, 15 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.13 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 1.24 (3 H, br m,  $\text{CH}_3\text{CH}_2$ -), 2.38 (2 H, m, piperidine  $\text{CH}$ -), 2.42 (2 H, m, piperidine  $\text{CH}$ -), 2.56 (4 H, m, piperidine  $\text{CH}$ -), 3.26 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.54 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}$ -), 3.68 (2 H, s,  $\text{CH}_2\text{N}$ -), 5.64 (1 H, br, NH), 6.12 (1 H, br, NH), 7.14 (3 H, m, ArH), 7.31 (3 H, m, ArH), 7.35 (1 H, m, ArH), 7.38 (1 H, m, ArH), 7.56 (1 H, s, ArH), 7.65 (2 H, m, ArH), 8.58 (1 H, m, ArH).

**N,N-Diethyl-4-[1-(3-thiophene)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-**

**benzamide (5e).**

Method as for **5a** using **4** (196 mg, 0.5 mmol) and 3-thiophenecarboxaldehyde (112 mg, 1.0 mmol) provided **5e** (185 mg, 76 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.13 (3 H, br m,  $\text{CH}_3\text{CH}_2-$ ), 1.24 (3 H, br m,  $\text{CH}_3\text{CH}_2-$ ), 2.46 (4 H, m, piperidine  $\text{CH}_2$ ), 2.84 (4 H, m, piperidine  $\text{CH}_2$ ), 3.26 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}-$ ), 3.52 (2 H, br m,  $\text{CH}_3\text{CH}_2\text{N}-$ ), 3.91 (2 H, s,  $\text{CH}_2\text{N}-$ ), 5.72 (1 H, br, NH), 6.44 (1 H, br, NH), 7.13 (3 H, m, ArH), 7.26 (3 H, m, ArH), 7.36 (3 H, m, ArH), 7.61 (1 H, s, ArH), 7.68 (1 H, d,  $J = 7.6$  Hz, ArH); *Anal. Calcd. for*  $\text{C}_{29}\text{H}_{33}\text{N}_3\text{O}_2\text{S} \cdot 2.5 \text{ HCl}$ : C, 60.18 %; H, 6.18 %; *Found*: C, 60.16 %; H, 6.49 %.

**N,N-Diethyl-4-[1-(2-thiazole)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5f).**

To a solution of N,N-diethyl-4-[piperidin-4-ylidene(3-carbamoylphenyl)-methyl]-benzamide (**4**, 300 mg, 0.8 mmol) in 1,2-dichloroethane (15 mL) was added 2-thiazole carboxaldehyde (94  $\mu\text{L}$ , 1.1 mmol) and sodium triacetoxyborohydride (228 mg, 1.1 mmol). The reaction mixture was stirred for 20h at room temperature, and then quenched with aqueous  $\text{NaHCO}_3$ . The aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  (2 x 20 mL) and the combined organic phases were dried over  $\text{MgSO}_4$ , filtered and evaporated. The crude product was purified by flash chromatography to give product (**5f**) as a yellow foam (234 mg, 63% yield).

The product was dissolved in dichloromethane (5mL) and a solution of HCl in ether (1N, 1.4 mL, 3eq.) was added. After 30 minutes the suspension was concentrated and the solid dried.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.10 (3H, t,  $J=7\text{Hz}$ ,  $\text{CH}_3$ ) ; 1.21 (3H, t,  $J=7\text{Hz}$ ,  $\text{CH}_3$ ) ; 2.64 (4H, br s,  $\text{CH}_2$ ) ; 3.27-3.31 (4H, m,  $\text{CH}_2$ ) ; 3.48-3.53 (2H, m,  $\text{CH}_2$ ) ; 3.67 (2H, br s,  $\text{CH}_2$ ) ; 4.77 (2H, s,  $\text{NCH}_2\text{Ar}$ ) ; 7.27 (2H, d,  $J=8.5\text{Hz}$ , Ar-H) ; 7.33-7.37 (3H, m, Ar-H) ; 7.44 (1H, t,  $J=7.5\text{Hz}$ , Ar-H) ; 7.68-7.69 (1H, m, Ar-H) ; 7.75-7.78 (2H, m, Ar-H) ; 7.94 (1H, d,  $J=3.5\text{Hz}$ , Ar-H). *Anal. Calcd for*  $\text{C}_{28}\text{H}_{32}\text{N}_4\text{O}_2\text{S} \cdot 2.5\text{HCl}$ : C, 58.00%; H, 6.00%; N, 9.66%; *Found*: C, 58.00%; H, 5.95%; N, 9.43%.

**N,N-Diethyl-4-[1-(3-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5g).**

Method as for **5f** using **4** (176 mg, 0.45 mmol) and 3-pyridinecarboxyaldehyde (60  $\mu\text{L}$ , 0.6 mmol) provided **5g** (91.5 mg, 42 %):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.09 (3H, t,  $J=7\text{Hz}$ ,  $\text{CH}_3$ ) ; 1.21

(3H, t, J=7Hz, CH<sub>3</sub>) ; 2.68-2.75 (4H, m, CH<sub>2</sub>) ; 3.26-3.28 (4H, m, CH<sub>2</sub>) ; 3.43-3.60 (4H, m, CH<sub>2</sub>) ; 4.65 (2H, s, NCH<sub>2</sub>Ar) ; 7.27 (2H, d, J=8.5Hz, Ar-H) ; 7.32-7.36 (3H, m, Ar-H) ; 7.43 (1H, t, J=8Hz, Ar-H) ; 7.70-7.71 (1H, m, Ar-H) ; 7.75-7.78 (1H, m, Ar-H) ; 8.19 (1H, dd, J=6, 8Hz, Ar-H) ; 8.88 (1H, d, J=8Hz, Ar-H) ; 8.98 (1H, d, J=6Hz, Ar-H) ; 9.21 (1H, s, Ar-H). *Anal.* Calcd for C<sub>30</sub>H<sub>34</sub>N<sub>4</sub>O<sub>2</sub> x 3.1HCl x 0.4H<sub>2</sub>O: C, 59.77%; H, 6.34%; N, 9.29%; *Found*: C, 59.70%; H, 6.36%; N, 9.17%.

**N,N-Diethyl-4-[1-(2-pyrrole)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5h).**

Method as for **5f** using **4** (261 mg, 0.7 mmol) and 2-pyrrolecarboxyaldehyde (89 mg, 0.9 mmol) provided **5h** (118.5 mg, 38 %): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.09 (3H, t, J=7Hz, CH<sub>3</sub>) ; 1.21 (3H, t, J=7Hz, CH<sub>3</sub>) ; 2.46-2.53 (2H, m, CH<sub>2</sub>) ; 2.65-2.77 (2H, m, CH<sub>2</sub>) ; 2.97-3.04 (2H, m, CH<sub>2</sub>) ; 3.26-3.30 (2H, m, CH<sub>2</sub>) ; 3.46-3.52 (4H, m, CH<sub>2</sub>) ; 4.30 (2H, s, NCH<sub>2</sub>Ar) ; 6.33-6.34 (1H, m, Ar-H) ; 6.85-6.86 (1H, m, Ar-H) ; 7.24-7.26 (2H, m, Ar-H) ; 7.30-7.36 (4H, m, Ar-H) ; 7.40-7.45 (1H, m, Ar-H) ; 7.67-7.68 (1H, m, Ar-H) ; 7.75-7.77 (1H, m, Ar-H). *Anal.* Calcd for C<sub>29</sub>H<sub>34</sub>N<sub>4</sub>O<sub>2</sub> x 1.1HCl x 1.8H<sub>2</sub>O: C, 64.13%; H, 7.18%; N, 10.32%; *Found*: C, 64.26%; H, 7.15%; N, 9.94%.

**N,N-Diethyl-4-[1-(4-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5i).**

Method as for **5f** using **4** (329 mg, 0.8 mmol) and 4-pyridinecarboxyaldehyde (112 μL, 1.2 mmol) provided **5i** (217 mg, 54 %): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.12 (3H, t, J=7Hz, CH<sub>3</sub>) ; 1.24 (3H, t, J=7Hz, CH<sub>3</sub>) ; 2.67-2.82 (4H, m, CH<sub>2</sub>) ; 3.22-3.34 (4H, m, CH<sub>2</sub>) ; 3.49-3.65 (4H, m, CH<sub>2</sub>) ; 4.72 (2H, s, NCH<sub>2</sub>Ar) ; 7.29 (2H, d, J=8.5Hz, Ar-H) ; 7.33-7.40 (3H, m, Ar-H) ; 7.46 (1H, t, J=8Hz, Ar-H) ; 7.72-7.73 (1H, m, Ar-H) ; 7.78-7.80 (1H, m, Ar-H) ; 8.37 (2H, d, J=7Hz, Ar-H) ; 9.00 (2H, d, J=7Hz, Ar-H).

**N,N-Diethyl-4-[1-(4-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide (5j).**

Method as for **5f** using **4** (313 mg, 0.8 mmol) and 4-imidazolecarboxyaldehyde (108 mg, 1.1 mmol) provided **5j** (68.2 mg, 18 %): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.12 (3H, t, J=7Hz, CH<sub>3</sub>) ; 1.23 (3H, t, J=7Hz, CH<sub>3</sub>) ; 2.63 (2H, t, J=6Hz, CH<sub>2</sub>) ; 2.68 (2H, t, J=6Hz, CH<sub>2</sub>) ; 3.26-3.36 (6H, m, CH<sub>2</sub>)



;3.54 (2H, q, J=7Hz, CH<sub>2</sub>) ; 4.45 (2H, s, NCH<sub>2</sub>Ar) ; 7.28 (2H, d, J=8Hz, Ar-H) ; 7.32-7.40 (3H, m, Ar-H) ; 7.45 (1H, t, J=8Hz, Ar-H) ; 7.66 (1H, s, Ar-H) ; 7.71 (1H, t, J=2Hz, Ar-H) ; 7.76-7.82 (1H, m, Ar-H) ; 8.61 (1H, s, Ar-H).

## 5 Pharmaceutical compositions

The novel compounds according to the present invention may be administered orally, intramuscularly, subcutaneously, topically, intranasally, intraperitoneally, intrathoracically, intravenously, epidurally, intrathecally, intracerebroventricularly and by injection into the joints.

A preferred route of administration is orally, intravenously or intramuscularly.

10 The dosage will depend on the route of administration, the severity of the disease, age and weight of the patient and other factors normally considered by the attending physician, when determining the individual regimen and dosage level as the most appropriate for a particular patient.

For preparing pharmaceutical compositions from the compounds of this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include  
15 powders, tablets, dispersible granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, or tablet disintegrating agents; it can also be an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided  
20 active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

For preparing suppository compositions, a low-melting wax such as a mixture of fatty acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient sized molds and  
25 allowed to cool and solidify.

Suitable carriers are magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

Salts include, but are not limited to, pharmaceutically acceptable salts. Examples of pharmaceutically acceptable salts within the scope of the present invention include: acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, bromide, calcium acetate, camsylate, carbonate, chloride, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, glucaptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, isethionate, lactate, lactobionate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, pamoate (embonate), pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate. Examples of pharmaceutically unacceptable salts within the scope of the present invention include: hydroiodide, perchlorate, and tetrafluoroborate.

Preferred pharmaceutically acceptable salts are the hydrochlorides, sulfates and bitartrates.

The hydrochloride and sulfate salts are particularly preferred.

The term composition is intended to include the formulation of the active component with encapsulating material as a carrier providing a capsule in which the active component (with or without other carriers) is surrounded by a carrier which is thus in association with it. Similarly, cachets are included. Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

Liquid from compositions include solutions, suspensions, and emulsions. Sterile water or water-propylene glycol solutions of the active compounds may be mentioned as an example of liquid preparations suitable for parenteral administration. Liquid compositions can also be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a viscous material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl cellulose, and other suspending agents known to the pharmaceutical formulation art.

Preferably the pharmaceutical composition is in unit dosage form. In such form, the composition is divided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of the preparations, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself, or it can be the appropriate number of any of these packaged forms.

## **BIOLOGICAL EVALUATION**

### **In vitro model**

#### **Cell culture**

A. Human 293S cells expressing cloned human  $\mu$ ,  $\delta$ , and  $\kappa$  receptors and neomycin resistance were grown in suspension at 37°C and 5% CO<sub>2</sub> in shaker flasks containing calcium-free DMEM10% FBS, 5% BCS, 0.1% Pluronic F-68; and 600  $\mu$ g/ml geneticin.

B. Mouse and rat brains were weighed and rinsed in ice-cold PBS (containing 2.5mM EDTA, pH 7.4). The brains were homogenized with a polytron for 15 sec (mouse) or 30 sec (rat) in ice-cold lysis buffer (50mM Tris, pH 7.0, 2.5mM EDTA, with phenylmethylsulfonyl fluoride. added just prior use to 0.5mM from a 0.5M stock in DMSO:ethanol).

#### **Membrane preparation**

Cells were pelleted and resuspended in lysis buffer (50 mM Tris, pH 7.0, 2.5 mM EDTA, with PMSF added just prior to use to 0.1 mM from a 0.1 M stock in ethanol), incubated on ice for 15 min, then homogenized with a polytron for 30 sec. The suspension was spun at 1000 g (max) for 10 min at 4 °C. The supernatant was saved on ice and the pellets resuspended and spun as before. The supernatants from both spins were combined and spun at 46,000 g (max) for 30 min. The pellets were resuspended in cold Tris buffer (50 mM Tris/Cl, pH 7.0) and spun again. The final pellets were resuspended in membrane buffer (50 mM Tris, 0.32 M sucrose, pH 7.0). Aliquots (1 ml) in polypropylene tubes were frozen in dry ice/ethanol and stored at -70 °C until use. The protein concentrations were determined by a modified Lowry assay with sodium dodecyl sulfate.



### **Binding assays**

Membranes were thawed at 37 °C, cooled on ice, passed 3 times through a 25-gauge needle, and diluted into binding buffer (50 mM Tris, 3 mM MgCl<sub>2</sub>, 1 mg/ml BSA (Sigma A-7888), pH 7.4, which was stored at 4°C after filtration through a 0.22 m filter, and to which had been  
5 freshly added 5 µg/ml aprotinin, 10 µM bestatin, 10 µM diprotin A, no DTT). Aliquots of 100 µl were added to iced 12x75 mm polypropylene tubes containing 100 µl of the appropriate radioligand and 100 µl of test compound at various concentrations. Total (TB) and nonspecific (NS) binding were determined in the absence and presence of 10 µM naloxone respectively. The tubes were vortexed and incubated at 25°C for 60-75 min, after which time the contents are rapidly vacuum-  
10 filtered and washed with about 12 ml/tube iced wash buffer (50 mM Tris, pH 7.0, 3 mM MgCl<sub>2</sub>) through GF/B filters (Whatman) presoaked for at least 2h in 0.1% polyethyleneimine. The radioactivity (dpm) retained on the filters was measured with a beta counter after soaking the filters for at least 12h in minivials containing 6-7 ml scintillation fluid. If the assay is set up in 96-place deep well plates, the filtration is over 96-place PEI-soaked unifilters, which were washed with 3 x 1 ml  
15 wash buffer, and dried in an oven at 55°C for 2h. The filter plates were counted in a TopCount (Packard) after adding 50 µl MS-20 scintillation fluid/well.

### **Functional Assays**

The agonist activity of the compounds is measured by determining the degree to which the compounds receptor complex activates the binding of GTP to G-proteins to which the receptors are  
20 coupled. In the GTP binding assay, GTP[γ]<sup>35</sup>S is combined with test compounds and membranes from HEK-293S cells expressing the cloned human opioid receptors or from homogenised rat and mouse brain. Agonists stimulate GTP[γ]<sup>35</sup>S binding in these membranes. The EC<sub>50</sub> and E<sub>max</sub> values of compounds are determined from dose-response curves. Right shifts of the dose response curve by the delta antagonist naltrindole are performed to verify that agonist activity is mediated through  
25 delta receptors.




### **Data analysis**




The specific binding (SB) was calculated as TB-NS, and the SB in the presence of various test compounds was expressed as percentage of control SB. Values of IC<sub>50</sub> and Hill coefficient (n<sub>H</sub>) for ligands in displacing specifically bound radioligand were calculated from logit plots or curve fitting

programs such as Ligand, GraphPad Prism, SigmaPlot, or ReceptorFit. Values of  $K_i$  were calculated from the Cheng-Prussoff equation. Mean  $\pm$  S.E.M. values of  $IC_{50}$ ,  $K_i$  and  $n_H$  were reported for ligands tested in at least three displacement curves. Biological data are tabulated on the following pages in Table 1.

5

Table 1: Biological data.

Ex. #	MOLECULAR STRUCTURE	HDELTA (nM)			RAT BRAIN		MOUSE BRAIN	
		IC <sub>50</sub>	EC <sub>50</sub>	%EMax	EC <sub>50</sub>	%EMax	EC <sub>50</sub>	%EMax
3b		0.548	0.091	93.775	0.403	178.94	0.539	164.82
5a		0.373	0.158	98.107	0.613	181.01	0.818	170.24
5b		0.282			0.633	149.28		

<b>5c</b>		0.246			0.367	170.4		
<b>5d</b>		0.278			0.396	179.89		
<b>5e</b>		0.235						

### Receptor saturation experiments

Radioligand  $K_d$  values were determined by performing the binding assays on cell membranes with the appropriate radioligands at concentrations ranging from 0.2 to 5 times the estimated  $K_d$  (up to 10 times if amounts of radioligand required are feasible). The specific radioligand binding was expressed as pmole/mg membrane protein. Values of  $K_d$  and  $B_{max}$  from individual experiments were obtained from nonlinear fits of specifically bound (B) vs. nM free (F) radioligand from individual according to a one-site model.

### Determination Of Mechano-Allodynia Using Von Frey Testing

Testing was performed between 08:00 and 16:00h using the method described by Chaplan et al. (1994). Rats were placed in Plexiglas cages on top of a wire mesh bottom which allowed access to the paw, and were left to habituate for 10-15 min. The area tested was the mid-plantar

left hind paw, avoiding the less sensitive foot pads. The paw was touched with a series of 8 Von Frey hairs with logarithmically incremental stiffness (0.41, 0.69, 1.20, 2.04, 3.63, 5.50, 8.51, and 15.14 grams; Stoelting, Ill, USA). The von Frey hair was applied from underneath the mesh floor perpendicular to the plantar surface with sufficient force to cause a slight buckling against the paw, and held for approximately 6-8 seconds. A positive response was noted if the paw was sharply withdrawn. Flinching immediately upon removal of the hair was also considered a positive response. Ambulation was considered an ambiguous response, and in such cases the stimulus was repeated.

### **Testing Protocol**

The animals were tested on postoperative day 1 for the FCA-treated group. The 50% withdrawal threshold was determined using the up-down method of Dixon (1980). Testing was started with the 2.04 g hair, in the middle of the series. Stimuli were always presented in a consecutive way, whether ascending or descending. In the absence of a paw withdrawal response to the initially selected hair, a stronger stimulus was presented; in the event of paw withdrawal, the next weaker stimulus was chosen. Optimal threshold calculation by this method requires 6 responses in the immediate vicinity of the 50% threshold, and counting of these 6 responses began when the first change in response occurred, e.g. the threshold was first crossed. In cases where thresholds fell outside the range of stimuli, values of 15.14 (normal sensitivity) or 0.41 (maximally allodynic) were respectively assigned. The resulting pattern of positive and negative responses was tabulated using the convention, X = no withdrawal; O = withdrawal, and the 50% withdrawal threshold was interpolated using the formula:

$$50\% \text{ g threshold} = 10^{(X_f + k\delta)} / 10,000$$

where  $X_f$  = value of the last von Frey hair used (log units);  $k$  = tabular value (from Chaplan et al. (1994)) for the pattern of positive / negative responses; and  $\delta$  = mean difference between stimuli (log units). Here  $\delta = 0.224$ .

Von Frey thresholds were converted to percent of maximum possible effect (% MPE), according to Chaplan et al. 1994. The following equation was used to compute % MPE:

$$\% \text{MPE} = \frac{\text{Drug treated threshold (g)} - \text{allodynia threshold (g)} \times 100}{\text{Control threshold (g)} - \text{allodynia threshold (g)}}$$

### Administration of Test Substance

Rats were injected (subcutaneously, intraperitoneally, intravenously or orally) with a test substance prior to von Frey testing, the time between administration of test compound and the von Frey test varied depending upon the nature of the test compound.

### 5 Writhing Test

Acetic acid will bring abdominal contractions when administered intraperitoneally in mice. These will then extend their body in a typical pattern. When analgesic drugs are administered, this described movement is less frequently observed and the drug selected as a potential good candidate.

A complete and typical Writhing reflex is considered only when the following elements are  
10 present: the animal is not in movement; the lower back is slightly depressed; the plantar aspect of *both* paws is observable. In this assay, compounds of the present invention demonstrate significant inhibition of writhing responses after oral dosing of 1-100  $\mu\text{mol/kg}$ .

#### **(i) Solutions preparation**

Acetic acid (AcOH): 120  $\mu\text{L}$  of Acetic Acid is added to 19.88 ml of distilled water in  
15 order to obtain a final volume of 20 ml with a final concentration of 0.6% AcOH. The solution is then mixed (vortex) and ready for injection.

Compound (drug): Each compound is prepared and dissolved in the most suitable vehicle according to standard procedures.

#### **(ii) Solutions administration**

20 The compound (drug) is administered orally, intraperitoneally (i.p.), subcutaneously (s.c.) or intravenously (i.v.) at 10 ml/kg (considering the average mice body weight) 20, 30 or 40 minutes (according to the class of compound and its characteristics) prior to testing. When the compound is delivered centrally: Intraventricularly (i.c.v.) or intrathecally (i.t.) a volume of 5  $\mu\text{L}$  is administered.

The AcOH is administered intraperitoneally (i.p.) in two sites at 10 ml/kg (considering the  
25 average mice body weight) immediately prior to testing.

#### **(iii) Testing**


The animal (mouse) is observed for a period of 20 minutes and the number of occasions (Writhing reflex) noted and compiled at the end of the experiment. Mice are kept in individual "shoe box" cages with contact bedding. A total of 4 mice are usually observed at the same time: one control and three doses of drug.

5

10





CLAIMS


1. A compound of the formula I




wherein

$R^1$  is selected from any one of



(vii) pyrrolyl  ; and


(viii) thiazolyl  ;


5 where each  $R^1$  phenyl ring and  $R^1$  heteroaromatic ring may independently be further substituted by 1, 2 or 3 substituents independently selected from straight and branched  $C_1$ - $C_6$  alkyl,  $NO_2$ ,  $CF_3$ ,  $C_1$ - $C_6$  alkoxy, chloro, fluoro, bromo, and iodo; as well as salts thereof.

10 2. A compound according to claim 1 wherein  $R^1$  is selected from any one of

(i) phenyl  ;

(ii) pyridinyl  ;

15 (iii) thienyl  ; and

(iv) furanyl  ;

20 3. A compound according to claim 1, wherein each  $R^1$  phenyl ring and  $R^1$  heteroaromatic ring may optionally and independently be further substituted by 1, 2 or 3 substituents independently selected from methyl,  $CF_3$ , chloro, fluoro, bromo, and iodo.

25 4. A compound according to claim 1, selected from any one of

N,N-Diethyl-4-[piperidin-4-ylidene(3-carbamoylphenyl)-methyl]-benzamide;



N,N-Diethyl-4-[1-(2-thiophene)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

N,N-Diethyl-4-[1-(2-furfuryl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl)-benzamide;

N,N-Diethyl-4-[1-(3-furfuryl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl)-benzamide;

5 N,N-Diethyl-4-[1-(2-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

N,N-Diethyl-4-[1-(3-thiophene)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

N,N-Diethyl-4-[1-(2-thiazole)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

N,N-Diethyl-4-[1-(3-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

10 N,N-Diethyl-4-[1-(2-pyrrole)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;

N,N-Diethyl-4-[1-(4-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide;  
and

N,N-Diethyl-4-[1-(4-pyridine)methyl-piperidin-4-ylidene-(3-carbamoylphenyl)-methyl]-benzamide.

15 5. A compound according to Claim 1, in form of its hydrochloride, dihydrochloride, sulfate, tartrate, ditrifluoroacetate or citrate salts.

6. A compound according to any one of claims 1-5 for use in therapy.

20 7. A compound according to claim 6, wherein the therapy is pain management.

8. A compound according to claim 6, wherein the therapy is directed towards gastrointestinal disorders.

25 9. A compound according to claim 6, wherein the therapy is directed towards spinal injuries.

10. A compound according to claim 6, wherein the therapy is directed to disorders of the sympathetic nervous system.

30 11. Use of a compound according to formula I of claim 1 for the manufacture of a medicament for use in the treatment of pain.

12. Use of a compound according to formula I of claim 1 for the manufacture of a medicament for use in the treatment of gastrointestinal disorders.

5 13. Use of a compound according to formula I of claim 1 for the manufacture of a medicament for use in the treatment of spinal injuries.

14. A pharmaceutical composition comprising a compound of the formula I according to claim 1 as an active ingredient, together with a pharmaceutically acceptable carrier.

10 15. A method for the treatment of pain, whereby an effective amount of a compound of the formula I according to claim 1 is administered to a subject in need of pain management.

16. A method for the treatment of gastrointestinal disorders, whereby an effective amount of a  
15 compound of the formula I according to claim 1, is administered to a subject suffering from said gastrointestinal disorder.

17. A method for the treatment of spinal injuries, whereby an effective amount of a compound of the formula I according to claim 1, is administered to a subject suffering from said spinal injury.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 02/01804

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 211/70, C07D 401/06, C07D 407/06, C07D 409/06, C07D 417/06,  
A61K 31/4523, A61K 31/445, A61P 25/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CHEM. ABS DATA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9828275 A1 (ASTRA PHARMA INC.), 2 July 1998 (02.07.98), the claims, esp. claims 1-3, 11-14 and claim 4, page 112 --	1-17
P,X	WO 0174804 A1 (ASTRAZENECA AB), 11 October 2001 (11.10.01) -- -----	1-17

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier application or patent but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 "&" document member of the same patent family

Date of the actual completion of the international search

24 January 2003

Date of mailing of the international search report

30-01-2003

Name and mailing address of the ISA/  
 Swedish Patent Office  
 Box 5055, S-102 42 STOCKHOLM  
 Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson/Eö  
 Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE02/01804

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 15-17  
because they relate to subject matter not required to be searched by this Authority, namely:  
**see next sheet**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE02/01804

Claims 15-17 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 02/01804

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9828275	A1	02/07/98	AU	737999 B	06/09/01
				AU	5351298 A	17/07/98
				BR	9714055 A	09/05/00
				CN	1246111 A	01/03/00
				CZ	9902199 A	17/11/99
				EE	3824 B	15/08/02
				EE	9900256 A	15/12/99
				EP	0946511 A	06/10/99
				IL	130535 D	00/00/00
				JP	2001507350 T	05/06/01
				NO	313670 B	11/11/02
				NO	993022 A	20/08/99
				NZ	336029 A	30/03/01
				PL	334374 A	28/02/00
				SE	9604785 D	00/00/00
				SK	76299 A	08/11/99
				TR	9901417 T	00/00/00
				US	6187792 B	13/02/01
				US	6455545 B	24/09/02
				US	2001021715 A	13/09/01
				ZA	9711059 A	22/06/98
				HU	0000610 A	28/09/00
				SE	9702535 D	00/00/00
WO	0174804	A1	11/10/01	AU	4498101 A	15/10/01
				AU	5584100 A	02/01/01
				EP	1210277 A	05/06/02
				NO	20024775 D	00/00/00
				SE	0001207 D	00/00/00
				SK	17892001 A	09/05/02